

Croat Med J. 2014;55:156-62
doi: 10.3325/cmj.2014.55.156

Alcohol abuse as the strongest risk factor for violent offending in patients with paranoid schizophrenia

Aim To determine predictive risk factors for violent offending in patients with paranoid schizophrenia in Croatia.

Method The cross-sectional study including male in-patients with paranoid schizophrenia with (N=104) and without (N=102) history of physical violence and violent offending was conducted simultaneously in several hospitals in Croatia during one-year period (2010-2011). Data on their sociodemographic characteristics, duration of untreated illness phase (DUP), alcohol abuse, suicidal behavior, personality features, and insight into illness were collected and compared between groups. Binary logistic regression model was used to determine the predictors of violent offending.

Results Predictors of violent offending were older age, DUP before first contact with psychiatric services, and alcohol abuse. Regression model showed that the strongest positive predictive factor was harmful alcohol use, as determined by AUDIT test (odds ratio 37.01; 95% confidence interval 5.20-263.24). Psychopathy, emotional stability, and conscientiousness were significant positive predictive factors, while extroversion, pleasantness, and intellect were significant negative predictive factors for violent offending.

Conclusion This study found an association between alcohol abuse and the risk for violent offending in paranoid schizophrenia. We hope that this finding will help improve public and mental health prevention strategies in this vulnerable patient group.

Marija Kudumija
Slijepčević^{1,2}, Vlado
Jukić³, Darko Novalić¹,
Tija Žarković-Palijan⁴,
Milan Milošević⁵, Ivana
Rosenzweig^{6,7}

¹Department of Psychiatry, General Hospital Bjelovar, Bjelovar, Croatia

²Technical College Bjelovar, Bjelovar, Croatia

³Psychiatric Clinical Hospital Vrapče, Zagreb, Croatia

⁴Psychiatric Hospital "Dr Ivan Barbot", Popovača, Croatia

⁵Department for Environmental and Occupational Health, University of Zagreb, School of Medicine, Andrija Štampar School of Public Health, Zagreb, Croatia

⁶Department of Neuroimaging, Institute of Psychiatry, King's College London, London, United Kingdom

⁷Danish Epilepsy Centre, Dianalund, Denmark

Received: October 29, 2012
Accepted: February 13, 2014

Correspondence to:

Marija Kudumija Slijepčević
Department of Psychiatry
General Hospital Bjelovar
43000 Bjelovar, Croatia
mkudumija@gmail.com

Individuals with schizophrenia have an increased risk of violence (1), but different studies report different risks (1,2). Anglo-American studies commonly report higher prevalence rates than European studies (3,4). These patients have also been reported to have up to 4-6 times higher violent behavior rate than the general population (3-5). Nonetheless, less than 0.2% patients suffering from schizophrenia commit homicide (in 20-year period) and less than 10% of commit a violent act (3). Also, patients with schizophrenia contribute to 6%-11% of all homicides and homicide attempts (3-5).

In general, aggressiveness is usually associated with anti-social personality features, juvenile delinquency, and psychoactive substance abuse (6). In patients with schizophrenia violence and violent offending is associated with a great number of risk factors, such as premorbid affinity to violent behavior, alcohol abuse, younger age, lower socioeconomic status (6,7), deinstitutionalization, longer duration of untreated psychosis, later onset of first episode of psychosis (1,4,8), lower social status, broken families, asocial behavior of parents, loss of father at an early age, a new marriage partner in the family, and growing up in an orphanage (9).

Several studies (10-12) looked at four basic personality dimensions and their role in violence in patients with schizophrenic illness spectrum: impulse control, affect regulation, narcissism, and paranoid cognition. Impulsivity and immature affect regulation were associated with most neuropsychiatric disorders, and were particularly predictive of affinity for addictive disorders, while paranoid cognition and narcissism were predictive of violence acts (10-12).

The causes of schizophrenia may be genetic, early environmental, and epigenetic risk factors (13,14), which may further modulate the risk of violent offending among individuals with this disease (1,15). Until recently, very little has been reported about the predictive factors of violence and violent offending in the patient population in Croatia. The Croatian population has during the last two decades been exposed to environmental and socio-demographic changes (eg, Croatian War 1991-1995 and post-war period), which might have had an impact on predictive risk factors. Therefore, we conducted a cross-sectional study of in-patients with paranoid schizophrenia with or without history of physical violence and violent offending (inclusive of homicide) in several hospitals in Croatia during one-year period.

PATIENTS AND METHODS

Patients and their sociodemographics

The study took place between December 1, 2010 and December 1, 2011. It included 206 male adult inpatients aged between 18 and 60 years who met the criteria for paranoid schizophrenia according to the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) (16) and who agreed to be part of the study. Non-inclusion criteria included neurological or significant somatic illnesses, learning disability, or organic brain damage.

Of these patients, 102 (non-violent offending) were voluntary inpatients at two acute psychiatric departments (General Hospital Bjelovar and Psychiatric Clinical Hospital Vrapče) with no recorded or reported history of violent offending. Open Aggressiveness Scale (OAS) (17) was administered at the admission and the cut-off point of 7 was established as the inclusion criterion (18).

Hundred and four male adult patients (violent offending group) were involuntary inpatients, hospitalized at forensic departments (Psychiatric Clinical Hospital Vrapče and Psychiatric Hospital "Dr Ivan Barbot" in Popovača) following the incidence of violence and violent offending, inclusive of homicide or attempted homicide. Twenty four patients left the study during assessments: 6 from non-violent offending group – due to inability to concentrate and 18 from violent offending group – 3 due to inability to concentrate, 4 due to sight impairment, 3 due to the fear that the results would be used against them, and 8 offered no explanation. Written informed consent was obtained from all participants and research was approved by the respective institutional ethics committees.

Methods

To insure consistency and intra- and inter-rater reliability, all the clinical assessments, data collection, and questionnaire application were carried out by a group of researchers and specialized psychiatrists. They were all trained for the purposes of this study and were tested for consistency for the used scales and DSM-IV criteria application for the diagnosis of paranoid schizophrenia (16).

Clinical assessment involved the following: structured psychiatric interview, collection of socio-demographic data (using a semi-structured questionnaire), duration of untreated psychosis (DUP) assessment (objective

and subjective), AUDIT questionnaire application (19), and InterSePT questionnaire application (20).

The following general and socio-demographic data were collected: age, age of the primary contact with psychiatric services, early psychomotor development history, family cohesion, parents' education, patient's educational, social, and employment status (inclusive of salary/pension level), marital status and the number of children, urbanicity of the place of residence and rural-urban migration, family psychiatric and medical history, and asocial behavior in the primary family.

For the assessment of alcohol consumption we used the AUDIT questionnaire, which covers three separable domains: consumption, harmful use, and dependent use (19). The score of 8 and more was used as the limit for determining harmful alcohol use (21). For the assessment of current

suicidal ideation we used 12-item InterSePTScale for Suicidal Thinking (20). DUP was assessed by clinical interview and the recorded psychiatric history in medical files. The presence of three groups of symptoms was determined – positive symptoms (hallucinations, delusions, and odd beliefs thought disorder), negative symptoms (depression, dysphoria, apathy, anergia, apathy, and amotivation), and signs of social decline (withdrawn behavior, poor interpersonal relationship, social avoidance, and lack of interest in education or work). The patients also filled in the International Personality Item Pool (IPIP, <http://ipip.ori.org/ipip/>) self-assessment questionnaires, Brief Cognitive Insight Scale (BCIS) (22), and they self-assessed how mentally unwell they had been before their first psychiatric treatment (DUP self-assessment). Psychopathy was defined as the sum score of all IPIP items that described sociability, assertiveness, emotional reactivity, irritability, impoliteness, need to control, and irresponsibility (23).

TABLE 1. General patients' characteristics (χ^2 test)

	Group				P	
	non-violent offending, N = 96		violent offending, N = 86			
	No.	%	No.	%		
Education	without school	0	0.00	2	2.33	<0.001
	elementary	17	17.71	25	29.07	
	high school	66	68.75	53	61.63	
	college	0	0.00	6	6.98	
	university	13	13.54	0	0.00	
Marriage status	no	75	78.13	76	93.83	0.003
	yes	21	21.88	5	6.17	
Significant somatic co-morbidity and/or history	no	81	84.38	71	82.56	0.742
	yes	15	15.63	15	17.44	
Head injury	no	63	65.63	45	52.94	0.083
	yes	33	34.38	40	47.06	
Urbanicity	no	26	27.08	48	56.47	<0.001
	yes	70	72.92	37	43.53	
Divorced parents	no	91	94.79	71	84.52	0.022
	yes	5	5.21	13	15.48	
Problems with law	no	76	79.17	75	88.24	0.102
	yes	20	20.83	10	11.76	
Monthly income (HRK)	<1000	30	34.09	37	44.05	0.001
	1000-3000	29	32.95	38	45.24	
	3000-5000	13	14.77	8	9.52	
	>5000	16	18.18	1	1.19	
Positive psychiatric family history	no	76	79.17	65	76.47	0.663
	yes	20	20.83	20	23.53	
Age, median (IQR)	34.00 (26.00-43.75)		43.50 (36.00-52.25)		<0.001*	
Duration of untreated psychosis (months), median (IQR)	8.00 (3.00-13.50)		12.00 (6.00-24.00)		<0.001*	
AUDIT score, median (IQR)	0.00 (0.00-4.00)		2.00 (0.00-8.00)		0.072*	

*Mann-Whitney U test.

Statistical analysis

Standard descriptive statistics measures (mean, standard deviation [SD], medians with interquartile ranges) were used. Normality of distribution was tested by Smirnov-Kolmogorov test, and suitable nonparametric methods were applied. Differences in quantitative values between the two groups were tested by a nonparametric Mann-Whitney U test, while differences in categorical variables were tested by χ^2 test. Binary logistic regression was performed to assess the impact of several predictor variables (age, education, AUDIT score, InterSept score, Beck's Cognitive Insight scale, psychopathy index, extraversion, agreeableness, consciousness, intellect, self-reflection, and duration of untreated psychosis) on the likelihood that patients had aggressive behavior. *P* values lower than or equal to 0.05 were considered significant. STATISTICA version 9.1 was used (www.statsoft.com).

RESULTS

General characteristics

Violent offending group was significantly older (43.80 years vs 34.83 years), but there was no significant difference in age of first contact with psychiatric services (violent offending = 27.69 years, non-violent offending = 24.23 years, $P > 0.05$, χ^2 test). Violent offending participants were five times more frequently children of divorced parents (84.5% vs 15.5%, $P = 0.022$, χ^2 test) and

significantly fewer of them had finished high school (13.5% non-violent offending vs 0% violent offending participants $P < 0.001$). Although there were no significant differences in school grades (the most common grades were good and very good in both groups, over 75% of participants), significantly more violent offending participants lost a year ($P = 0.001$, χ^2 test). Non-violent offending participants had significantly higher incomes (18.2% vs 1.2%, $P < 0.001$, χ^2 test) and were four times more likely to be married (Table 1).

Duration of the untreated illness phase

Both groups showed distorted insight as to when their psychosis had started and how long it had lasted (DUP self-assessment). The mean self-assessed DUP was 7.16 months in violent offending participants (SD 12.08; median 0.00 months) and 6.44 months in non-violent offending participants (SD 7.05, median 3.00 months, $P = 0.024$, Mann-Whitney test). Violent offending participants had significantly longer period of psychiatric assessment – 14.33 months (SD 9.20; median 12.00 months) than non-violent offending participants (9.45 months, SD 6.95; median 8.00 months, $P < 0.001$, Mann-Whitney Test).

Suicidal behavior, personality traits, insight into the illness, and alcohol abuse

No significant difference between the groups was found in suicidal thoughts and behavior ($P = 0.176$, Mann-Whitney

TABLE 2. Differences in individual questionnaire items and Beck's Cognitive Insight Scale (BCIS) subscales between violent offending and non-violent offending group (Mann-Whitney U-test)

	Group	N	Min	Max	Percentiles			<i>P</i>
					25th	Median	75th	
IPIP 50: Extraversion	violent offending	86	5.0	48.0	18.00	24.00	35.00	0.001
	non-violent offending	96	15.0	48.0	26.00	33.00	38.00	
IPIP 50: Agreeableness	violent offending	86	4.0	62.0	30.75	36.00	41.00	0.097
	non-violent offending	96	24.0	49.0	33.25	38.00	41.75	
IPIP 50: Consciousness	violent offending	86	3.0	50.0	36.00	39.00	44.00	0.038
	non-violent offending	96	19.0	48.0	34.00	37.00	41.00	
IPIP 50: Emotional stability	violent offending	86	3.0	50.0	27.75	34.50	40.25	0.011
	non-violent offending	96	16.0	44.0	26.00	29.00	37.00	
IPIP 50: Intellect	violent offending	86	3.0	50.0	26.00	33.00	38.25	0.111
	non-violent offending	96	19.0	50.0	31.00	34.00	39.75	
Self-reflectiveness subscale	violent offending	86	0.00	26.00	9.00	13.00	16.00	0.396
	non-violent offending	96	0.00	23.00	11.00	14.00	14.75	
Self-certainty subscale	violent offending	86	0.00	18.00	6.00	10.00	13.25	0.377
	non-violent offending	96	0.00	18.00	7.00	9.50	13.00	
BCIS composite index	violent offending	86	-12.00	15.00	0.00	2.50	6.00	0.218
	non-violent offending	96	-4.00	15.00	1.00	4.00	7.00	

test). InterSePT scale (20) had a satisfactory internal consistency (Cronbach $\alpha=0.82$). Non-violent offending group had a significantly higher extroversion score – 31.58 (SD 7.47; median 15.00) than violent offending group (27.093, SD 10.73; median 24.00, $P=0.001$, Mann-Whitney U test) (Table 2). Violent offending participants had a significantly higher mean conscientiousness score – 38.79 (SD 6.88; median 39.00, $P=0.038$, Mann-Whitney U test) and significantly higher mean emotional stability score – 33.21 (SD 9.72; median 34.50, $P=0.011$, Mann-Whitney U test). No significant differences were found between the groups in psychopathy index ($P=0.110$, Mann-Whitney U test) and in insight into the illness (Table 2). BCIS scale had a satisfactory internal consistency (Cronbach $\alpha=0.84$). Violent offending group was almost nine times more likely to be addicted to alcohol (AUDIT score >8 ; 27.1% vs 3.1%, $P<0.001$, χ^2 test).

Predictors for belonging to the violent-offending

Binary logistic regression model was significant (X^2_{13} test = 108.7, $P<0.001$). It explained 61.3% of variance and correctly classified 85.0% of participants. Hosmer and Lemeshow test was not significant ($P=0.201$). These data additionally indicate the validity of used regression model (Table 3).

The strongest positive violent offending predictor was AUDIT score higher than 8 (odds ratio [OR] 37.01, 95% confidence interval [CI] 5.20-263.24). These results suggest that the participants with the AUDIT score higher than 8 would have 37 times greater chances of belonging to violent offending group, with all other model variables controlled.

TABLE 3. Predictors of belonging to the violent offending group: binary logistic regression

	Odds 95% confidence		
	ratio	interval	P
Age	1.08	1.03-1.12	0.001
Education	0.60	0.30-1.19	0.142
AUDIT >8	37.01	5.20-263.24	<0.001
Clinical impression (InterSePT)	0.92	0.46-1.85	0.807
Psychopathy index	6.23	1.16-33.32	0.033
Extroversion	0.90	0.83-0.97	0.009
Agreeableness	0.89	0.81-0.98	0.019
Consciousness	1.13	1.02-1.25	0.016
Emotional stability	1.19	1.09-1.29	<0.001
Intellect	0.90	0.82-0.99	0.035
Beck's Cognitive Insight Scale	0.97	0.85-1.11	0.669
Self-reflectiveness	1.06	0.92-1.21	0.432
Duration of untreated psychosis (months)	1.10	1.05-1.16	<0.001

Other significant positive violent offending predictors included psychopathy index, emotional stability, conscientiousness, DUP, and age. Significant negative violent offending predictors (ie, the ones that reduced the chances of belonging to the violent offending group) were extroversion, pleasantness, and intellect (Table 3).

DISCUSSION

Our study found that alcohol addiction was the strongest violent offending predictor, even when adjusted for other model variables. Similarly, in Scandinavia, in one of the most extensive analyses to date, it was shown that substance abuse had a strong mediating effect on the risk for violent offending in schizophrenia (1). Substance abuse is prevalent among patients with schizophrenia and is frequently established at first presentation (24). In forensic settings, it was found that 26% of patients with schizophrenia and comorbid substance abuse were violent offenders, compared to only 7% patients without substance abuse (25). The same authors showed that individuals with both schizophrenia and substance abuse were 25.2 times more likely to commit violent crimes than healthy individuals (25). Abuse has been shown to be strongly associated with treatment non-compliance, tardive dyskinesia, criminality, and suicide (26-29).

Significantly higher rates of criminal conviction and recidivism have been found in patients with a lack of insight at discharge (30). We, on the other hand, did not find significant differences between the groups in insight into the illness. One explanation for this discrepancy is that our study was conducted in illness remission. Also, using only the BCIS as self-assessment questionnaire could have further biased our results. Objective clinical assessment of DUP suggested that this indeed might have been the case. Namely, DUP was significantly longer in the violent offending group. It has been suggested that the inability to recognize illness symptoms before first treatment could be connected with mentalization inability (31,32) and that this is another probable aggression predictor in patients suffering from schizophrenia. However, researchers are still somewhat divided on this topic and some authors have not found any causal relationship between violent behavior and DUP (33), while others have reported that DUP was connected with a worse illness prognosis, increased suicide risk, and possibly serious violent behavior (34).

In agreement with other studies, we also found that patients with a higher psychopathy index score had great-

er chances of belonging to the violent offending group (35,36), when all other model variables were controlled, although the comparison between the two groups did not reach significance. Moreover, the violent offending group also had lower education and was more emotionally stable than non-violent offending group.

Current clinical guidelines recommend that violence risk in schizophrenia should be consistently assessed but existing approaches can be resource-intensive and detailed clinical assessments of violence risk for most patients might not always be possible, especially in economically challenged health services (2). The observed predictive factors in the Croatian sample are in agreement with other previously published studies (1-3,34,37). However, in contrast to the study by Singh et al (2), "older age" in our study was more prognostic for violence and violent offending patients were on average a decade older (43.80 years) than patients with no history of violent offending (34.83 years). Of note, there was no significant difference in age of the first contact with psychiatric services. Also, since we used the cross-sectional design, our results can only suggest an association rather than point to a clear causal relationship. In conclusion, we hope that our findings will help relegate resources toward prevention of alcohol abuse in this vulnerable patient population.

Acknowledgments The authors thank Ivana Jurković, MA, for technical help and help in translation of the manuscript.

Funding None.

Ethical approval was received from Ethical committees Psychiatric Hospital in Popovača, Psychiatric Hospital Vrapče, and General Hospital Bjelovar.

Declaration of authorship MKS conceived the idea for the study, wrote the study protocol, performed data collection and analyses, and wrote the first draft. VJ wrote the protocol, performed data collection, and supervised the project. DN performed data collection and critically revised the manuscript. TŽP performed data collection and critically revised the manuscript. MM performed statistical analysis and critically revised the manuscript. IR critically revised the manuscript.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- Fazel S, Lingstrom N, Hjern A, Grann M, Lichtenstein P. Schizophrenia, substance abuse, and violent crime. *JAMA*. 2009;301:2016-23. [Medline:19454640](#) [doi:10.1001/jama.2009.675](#)
- Singh JP, Grann M, Lichtenstein P, Langstrom N, Fazel S. A novel approach to determining violence risk in schizophrenia: developing a stepped strategy in 13,806 discharged patients. *PLoS ONE*. 2012;7:e31727. [Medline:22359622](#) [doi:10.1371/journal.pone.0031727](#)
- Tiihonen J. Specific major mental disorders and criminality: a 26-year prospective study of the 1966 Northern Finland birth cohort. *Am J Psychiatry*. 1997;154:840-5. [Medline:9167513](#)
- Wallace C. Criminal offending in schizophrenia over 25-year period marked by deinstitutionalization and increasing prevalence of comorbid substance use disorder. *Am J Psychiatry*. 2004;161:716-27. [Medline:15056519](#) [doi:10.1176/appi.ajp.161.4.716](#)
- Mullen PE. Community care and criminal offending in schizophrenia. *Lancet*. 2000;355:614-7. [Medline:10696982](#) [doi:10.1016/S0140-6736\(99\)05082-5](#)
- Mulvey EP, Shaw E, Lidz CW. Why use multiple sources in research on patient violence in the community. *Crim Behav Ment Health*. 1994;4:235-58.
- Monahan J. Clinical and Actuarial predictions of violence. In: *Modern scientific evidence: the law and science of expert testimony*. Vol. 1. St Paul, MN: West Publishing Co; 1997. p. 300-18.
- Grann M, Fazel S. Substance misuse and violent crime: Swedish population study. *BMJ*. 2004;328:1233-4. [Medline:15155501](#) [doi:10.1136/bmj.328.7450.1233](#)
- Stompe T. Family and social influences on offending in man with schizophrenia. *Aust N Z J Psychiatry*. 2006;40:554-60. [Medline:16756580](#) [doi:10.1080/j.1440-1614.2006.01838.x](#)
- Taylor PJ, Leese M, Williams D, Butwell M, Daly R, Larkin E. Mental disorder and violence. A special (high security) hospital study. *Br J Psychiatry*. 1998;172:218-26. [Medline:9614470](#) [doi:10.1192/bjp.172.3.218](#)
- Šendula-Jengić V, Bošković G. Violence and criminal recidivism during forensic treatment [in Croatian]. *Med Jadert*. 2002;32:5-16.
- Nestor PG. Mental disorder and violence: personality dimensions and clinical features. *Am J Psychiatry*. 2002;159:1973-8. [Medline:12450942](#) [doi:10.1176/appi.ajp.159.12.1973](#)
- Vukadinovic Z, Rosenzweig I. Abnormalities in thalamic neurophysiology in schizophrenia: could psychosis be a result of potassium channel dysfunction? *Neurosci Biobehav Rev*. 2012;36:960-8. [Medline:22138503](#) [doi:10.1016/j.neubiorev.2011.11.005](#)
- Rosenzweig I, Vukadinovic Z, Turner AJ, Catani M. Neuroconnectivity and valproic acid: The myelin hypothesis. *Neurosci Biobehav Rev*. 2012;36:1848-56. [Medline:22652270](#) [doi:10.1016/j.neubiorev.2012.05.006](#)
- Kumari V, Gudjonsson GH, Raghuvanshi S, Barkataki I, Taylor P, Sumich A, et al. Reduced thalamic volume in men with antisocial personality disorder or schizophrenia and a history of serious violence and childhood abuse. *Eur Psychiatry*. 2013;28:225-34. [Medline:22944337](#) [doi:10.1016/j.eurpsy.2012.03.002](#)
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (4th ed.)*. Washington, DC: American Psychiatric Association; 2000.
- Yudofsky SC, Silver J, Jackson W. The overt aggression scale for the

- objective rating of verbal and physical aggression. *Am J Psychiatry*. 1986;143:35-9. [Medline:3942284](#)
- 18 Fresan A, Apiquian R, de la Fuente-Sandoval C. Premorbid adjustment and violent behavior in schizophrenic patients. *Schizophr Res*. 2004;69:143-8. [Medline:15469187](#) [doi:10.1016/j.schres.2003.07.002](#)
- 19 Rist F, Glockner-Rist A, Demmel R. The Alcohol Use Disorders Identification Test revisited: Establishing its structure using nonlinear factor analysis and identifying subgroups of respondents using latent class factor analysis. *Drug Alcohol Depend*. 2009;100:71-82. [Medline:19026497](#) [doi:10.1016/j.drugalcdep.2008.09.008](#)
- 20 Lindenmayer JP, Czobor P, Alphas L, Nathan AM, Anand R, Islam Z, et al. The InterSePT scale for suicidal thinking reliability and validity. *Schizophr Res*. 2003;63:161-70. [Medline:12892870](#) [doi:10.1016/S0920-9964\(02\)00335-3](#)
- 21 Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption. II. *Addiction*. 1993;88:791-804. [Medline:8329970](#) [doi:10.1111/j.1360-0443.1993.tb02093.x](#)
- 22 Millon T, Simonsen E, Smith MB, Davis RD. *Psychopathy*. New York: The Guilford Press; 1998.
- 23 Beck AT, Baruch E, Balter JM, Steer RA, Warman DM. A new instrument for measuring insight: the Beck Cognitive Insight Scale. *Schizophr Res*. 2004;68:319-29. [Medline:15099613](#) [doi:10.1016/S0920-9964\(03\)00189-0](#)
- 24 Mullen PE. Facing up to unpalatable evidence for the sake of our patients. *PLoS Med*. 2009;6:e1000112. [Medline:19688035](#) [doi:10.1371/journal.pmed.1000112](#)
- 25 Rice ME, Harris T. Psychopathy, schizophrenia, alcohol abuse and violent recidivism. *Int J Law Psychiatry*. 1995;18:333-42. [Medline:7591401](#) [doi:10.1016/0160-2527\(95\)00015-A](#)
- 26 Bailey I, Maxwell S, Brandabur MM. Substance abuse as a risk factor for tardive dyskinesia: a retrospective analysis of 1,027 patients. *Psychopharmacol Bull*. 1997;32:177-81.
- 27 Mueser KT, Yarnold P, Stanley R, Swett C, Miles K, Hill D. Substance use disorder in hospitalized severely mentally ill psychiatric patients: prevalence, correlates, and subgroups. *Schizophr Bull*. 2000;26:179-92. [Medline:10755680](#) [doi:10.1093/oxfordjournals.schbul.a033438](#)
- 28 Sullivan EV, Rosenbloom MJ, Serventi KL, Deshmukh A, Pfefferbaum A. Effects of alcohol dependence comorbidity and antipsychotic medication on volumes of the thalamus and pons in schizophrenia. *Am J Psychiatry*. 2003;160:1110-6. [Medline:12777269](#) [doi:10.1176/appi.ajp.160.6.1110](#)
- 29 De Bellis MD, Narasimhan A, Thatcher DL, Keshavan MS, Soloff P, Clark DB. Prefrontal cortex, thalamus, and cerebellar volumes in adolescents and young adults with adolescent-onset alcohol use disorders and comorbid mental disorders. *Alcohol Clin Exp Res*. 2005;29:1590-600. [Medline:16205359](#) [doi:10.1097/01.alc.0000179368.87886.76](#)
- 30 Soyka M, Graz C, Bottlender R, Dirschedl P, Schoech H. Clinical correlates of later violence and criminal offences in schizophrenia. *Schizophr Res*. 2007;94:89-98. [Medline:17509834](#) [doi:10.1016/j.schres.2007.03.027](#)
- 31 Addy K, Shannon K, Brookfield K. Theory of mind function, motor empathy, emotional empathy and schizophrenia: A single case study. *J Forensic Psychiatry Psychol*. 2007;18:293-306. [doi:10.1080/09670870701292746](#)
- 32 Fullam R, Dolan M. Criminal and personality profile of patients with schizophrenia and psychopathy. *Pers Individ Dif*. 2006;40:1591-602. [doi:10.1016/j.paid.2006.01.003](#)
- 33 Foley SR, Browne S, Clarke M, Kinsella A, Larkin C, O'Callaghan E. Is violence at presentation by patients with first-episode psychosis associated with duration of untreated psychosis? *Soc Psychiatry Psychiatr Epidemiol*. 2007;42:606-10. [Medline:17598060](#) [doi:10.1007/s00127-007-0217-9](#)
- 34 Large MM, Nielsens O, Ryan CJ, Hayes R. Mental health laws that require dangerousness for involuntary admission may delay the initial treatment of schizophrenia. *Soc Psychiatry Psychiatr Epidemiol*. 2008;43:251-6. [Medline:18060340](#) [doi:10.1007/s00127-007-0287-8](#)
- 35 Tengström A, Grann M, Långström N, Kullgren G. Psychopathy (PCL-R) as a predictor of violent recidivism among criminal offenders with schizophrenia. *Law Hum Behav*. 2000;24:45-58. [Medline:10693318](#) [doi:10.1023/A:1005474719516](#)
- 36 Belli H, Ural C. The association between schizophrenia and violent or homicidal behaviour: the prevention and treatment of violent behaviour in these patients. *West Indian Med J*. 2012;61:538-43. [Medline:23441379](#) [doi:10.7727/wimj.2011.028](#)
- 37 Jones RM, Lichtenstein P, Grann M, Långström N, Fazel S. Alcohol use disorders in schizophrenia: A national cohort study of 12,653 patients. *J Clin Psychiatry*. 2011;72:775-9. [Medline:21733478](#) [doi:10.4088/JCP.10m06320](#)