



## Središnja medicinska knjižnica

**Sakan S., Bandić Pavlović D., Milošević M., Virag I., Martinović P.,  
Dobrić I., Davila S., Perić M. (2015) *Implementing the Surgical Apgar  
Score in patients with trauma hip fracture. Injury, 46 (Suppl. 6). pp.  
S61-6. ISSN 0020-1383***

<http://www.elsevier.com/locate/issn/00201383>

<http://www.sciencedirect.com/science/journal/00201383>

<http://dx.doi.org/10.1016/j.injury.2015.10.051>

<http://medlib.mef.hr/2775>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

# Implementing the Surgical Apgar Score in the trauma hip fracture patient population

Sanja Sakan<sup>1</sup>, Daniela Bandic Pavlovic<sup>1</sup>, Milan Milosevic<sup>2</sup>, Igor Virag<sup>1</sup>, Petar Martinovic<sup>1</sup>, Ivan Dobric<sup>3</sup>, Slavko Davila<sup>3</sup>, Mladen Peric<sup>1</sup>

<sup>1</sup>Department of Anaesthesiology, Reanimatology and Intensive care, University Hospital Centre Zagreb, Zagreb, Croatia

<sup>2</sup>School of Public Health "Andrija Stampar", School of Medicine University of Zagreb, Zagreb, Croatia

<sup>3</sup>Department of Surgery, University Hospital Centre Zagreb, Zagreb, Croatia

Correspondence to: *Sanja Sakan, MD, Department of Anaesthesiology, Reanimatology and Intensive care, University Hospital Centre Zagreb, Kispaticeva 12, 10000 Zagreb, Croatia.*

*Phone : +385 95 902 4995; fax: +385 1 2376 014*

*e-mail: [sanja.sakan@hotmail.com](mailto:sanja.sakan@hotmail.com)*

## **Abstract**

**Backgrounds:** Trauma hip fractures in elderly patients are associated with a high postoperative morbidity and mortality, also premature death for a long-term period of many years. This high and long-term mortality among these patients can be explained not only by the fracture itself, but also due to patient's preoperative poor condition and comorbidities, the influence of the stressors such as surgery and type of anaesthesia on patient's condition, and the development of the major complications as cardiac failure, pulmonary embolism, pneumonia, deep venous thrombosis and acute renal failure in the postoperative period. Thus, the Surgical Apgar Score (SAS) could be a valuable clinicmetric tool that helps us objectively risk stratify patients immediately after the surgery, and enables those patients with the higher risk to have not only postoperative ICU care and good management during the hospital stay, but also after the hospital discharge as well.

**Methods:** The SAS was calculated retrospectively from the handwritten anaesthesia records on the 43 trauma hip fracture patients treated operatively in the University Hospital Center Zagreb over a 1-year period. The primary endpoints were the 30-days major postoperative complications and mortality, length of the ICU and hospital stay, and 6-months major complications development. Statistical analysis was applied to compare SAS with the patients' perioperative variables.

**Results:** The SAS score  $\leq 4$  in the trauma hip fracture patients was a significant predictor for the 30-days major surgical complications with 80% specificity (95% CI: 0.587-0.864,  $p=0.0111$ ). However the surgical score was not significant in the prediction of the 30-days mortality (95% CI: 0.468-0.771,  $p=0.2238$ ) and 6-months mortality (95% CI: 0.497-0.795,  $p=0.3997$ ) as primary endpoints in the hip fracture surgery patients.

**Conclusion:** Validity of the SAS is that it reveals the riddle how intraoperative events affect postoperative outcomes. On behalf ability of computing the SAS score in the operative theatre immediate, reliable, real-time feedback information about patient's postoperative risk stratification is gained. Nevertheless, our study showed that every trauma hip fracture patients with the  $SAS \leq 4$  should go postoperatively to the ICU and should be under intensive surveillance during the hospital stay as well after the hospital discharge too.

**Keywords:** Anaesthesia; hip fracture ;intensive care; Surgical Apgar Score; surgical outcome; trauma

## **Introduction**

Hip fractures in elderly population are usually the consequence of the weightless trauma due to poor bone quality (1). Likewise they are the second capital cause of the hospitalization increasing the postoperative complications and mortality rate of these patients for a short-term (30-days) and mid-term (6-months) period as well (2,3). According to the studies the novel 10-point Surgical Apgar Score (SAS) has been considered as a good independent predictor of the major postoperative complications and mortality within a period of 30-days after surgery among many surgical subspecialties (4). It is an easy calculated, simple, objective, real-time parameter computed as a sum of the three vital intraoperative variables. The variables are estimated blood loss (EBL), lowest heart rate (HR) and lowest mean arterial pressure (MAP) driven from the intraoperative anesthesia records data calculated at the end of surgery (5). Hence, we conducted a preliminary retrospective research to perceive a utility and validation of the SAS in rating the surgical outcome, length of the ICU and hospital stay in a trauma vulnerable geriatric hip fracture patients.

## **Patients and Methods**

### **Patient selection**

All necessary data of the patients were collected from the medical records and University Hospital Centre Zagreb electronic medical database to analyse the SAS importance. Our research included patients older than 18 years submitted to the hip fracture surgery and admitted postoperatively to the ICU in the University Hospital Centre Zagreb between March 1, 2013, and May 31, 2014. Beside already mentioned inclusion criteria, both sex, elective, expedited and emergent hip fracture surgery, general or spinal anaesthesia technique, and written informed consent were also eligible for inclusion in the study. Hip fracture patients with incomplete data were excluded. The 30-days and 6-months follow-up of the included patients was attained from the University Hospital Centre Zagreb electronic medical database and a private phone call to obtain a 6-months survival. The preliminary retrospective research protocol was approved by the Human Research Ethical Committee of the University Hospital Centre Zagreb

### **Calculation of the SAS**

The SAS is computed as a sum of the 3 intraoperative variables collected during the surgery starting from incision-to-skin closure time to minimise the episode of the anaesthetics reaction during induction and intubation on the hypotension and lowering heart rate (6). The variables are EBL, lowest HR and lowest MAP obtained from the intraoperative handwritten anaesthesia records for the each patient included in the study (5). Each of these 3 variables

according to the measured values were assigned particular scoring points after the SAS table (Figure 1). These 3 intraoperative variables points sum is a total SAS value of the particular patient for the specific operation (5).

### **Patient's preoperative characteristics and postoperative outcomes**

Patient's preoperative variables were bundled in four groups by organ system. Pulmonary comorbidity was defined as pneumonia, mechanical ventilator dependency, and preexisting chronic obstructive pulmonary disease. Cardiovascular comorbidity included earlier myocardial infarction, angina pectoris, congestive heart disease, coronary revascularization, peripheral vascular disease and anamnesis of stroke and transient ischaemic attack. Renal comorbidity included history of acute or chronic renal disease. Coagulation comorbidity comprised hereditary and acquired coagulation disorder, as well as the use of anticoagulation agents such as warfarin, acetylsalicylic acid and clopidogrel. Other preoperative variables were age, gender, American Society of Anaesthesiologists (ASA) physical status (from 1 to 4), length of operative delay (7). The preoperative patient's laboratory data consisted of haemoglobin, thrombocyte, and prothrombin time level.

The primary endpoints were occurrence of major postoperative complications and death within the follow-up period of 30-days after the hip fracture surgery, length of the ICU and hospital stay. The major complications were defined as the development of the following: postoperative bleeding that required transfusion of 4 units or more of packed red blood cells within 72 hours, cardiac arrest, myocardial infarction, deep venous thrombosis, pulmonary embolism, stroke or transient ischaemic attack, unplanned intubation, mechanical ventilation for 48 hours or more, pneumonia, sepsis, septic shock, and acute renal failure (8,9). Since we are without National Surgical Quality Improvement Program we have defined major postoperative complications according to other researches with the National Surgical Quality Improvement Program and reviewing the medical records. Complications were assessed by review of handwritten medical records, laboratory data, radiology records and electronic medical database by two independent researchers and verified by cross-examination. Six months postoperative mortality rate was obtained by a phone call to the patient or his family member.

### **Statistical analysis**

We performed a univariate analysis examining the relationship between each preoperative and intraoperative variables in the database and the outcomes of the major complication or death. We analysed categorical variables using Pearson Chi-square test. Kruskal-Wallis test was performed to assess continuous data normality and due to its results, appropriate non-parametric tests have been used. Mann-Whitney test was used to compare the SAS with

preoperative and intraoperative variables. Spearman rank correlation coefficients were used to correlate SAS with all other continuous variables. Receiver operating characteristics curve for the SAS value  $\leq 4$  and 30-days major complications. Binary logistic regression model has been made to assess prediction to group within 30 days postoperative major complications. Model included 4 predictor variables (ASA score, SAS score, age and gender) and major postoperative complications group as a binary dependent variable (0=without major complications and 1=with complication). P value below 0.05 was regarded significant. Statistical software MedCalc for Windows, version 13.0 was used for all analyses (MedCalc Statistical Software version 13.0 (MedCalc Software bvba, Ostend, Belgium; <http://www.medcalc.org>; 2014)

## Results

Our study included 43 cases, 8 men and 35 women. Baseline demographic and clinical characteristics of the cases are listed in the Table 1 and 2. Major complications within the 30-days occurred in 8 cases, hospital mortality in 1 case, 30-days mortality in 5 cases, and major complications within 6-months in 4 cases (Table 1). The median of operative delay was 3 days (25%-75% interquartile range (IQR) 2-6) (Table 2). Mean surgical score was 5.53 ( $\pm 1.79$  SD), and the SAS median was 6 (IQR 4-7) (Table 2). When we examined the relationship between the surgical score and 30-days major complications we obtained the SAS cut-off value 4 according to the area under the receiver operating characteristic curve (ROC AUC). The SAS value  $\leq 4$  was a significant predictor for the 30-days major surgical complications with 80% specificity (ROC AUC 0.743, SE 0.0957, 95% CI: 0.587-0.864,  $p=0.0111$ ) (Figure 2). However the surgical score was not significant in the prediction of the 30-days mortality (ROC AUC 0.629, SE 0.106, 95% CI: 0.468-0.771,  $p=0.2238$ ) and 6-months mortality (ROC AUC 0.657, SE 0.186, 95% CI: 0.497-0.795,  $p=0.3997$ ) as primary endpoints in the hip fracture surgery patients. According to our research other parameter must be relevant predictor for the 30-days and 6-months mortality primary endpoints. When we compared variables of cases with the SAS  $\leq 4$  (N=12) and the SAS  $\geq 5$  (N=31) with Pearson's Chi-Square Test, significant difference ( $p<0.05$ ) was found in the following variables: intraoperative EBL ( $p=0.016$ ,  $df=3$ ), intraoperative vasoactive support with noradrenaline or ephedrine bolus doses ( $p=0.022$ ,  $df=1$ ), and 30-days major complications ( $p=0.016$ ,  $df=1$ ) (Table 3). The SAS  $\leq 4$  group had greater intraoperative EBL. In this group 5 cases lost  $>1000$  ml blood, 4 cases between 601-1000ml, 3 cases between 101-600ml versus the SAS  $\geq 5$  group where 3 cases lost  $>1000$  ml blood, 5 cases 601-1000ml, 14 cases 101-600ml, and 9 cases  $\leq 100$  ml (Table 3). In the SAS  $\leq 4$  group vasoactive support was applied in 4 cases versus the SAS  $\geq 5$  group in 2 cases (Table 3). The 30-days

major complications happened in the SAS $\leq$ 4 group in 5 cases compared to the SAS  $\geq$ 5 group in 3 cases (Table 3). When the SAS was compared with patients' variables and postoperative outcomes using nonparametric Mann-Whitney test where appropriate, the significant correlation was found between the SAS and the lowest HR( $p=0.007$ ) (Table 2). Spearman rank correlation coefficients were used to correlate the SAS value with patients' preoperative and intraoperative variables, length of the ICU and hospital stay. The significant negative correlation was observed with the intraoperative EBL (the lower intraoperative bleeding, higher SAS,  $p<0.001$ ) and the lowest HR( lower HR, higher SAS,  $p<0.001$ ) (Table 4). Interestingly, this test observed that duration of surgery was also in negative correlation with the SAS (longer duration of surgery, lower SAS,  $p=0.014$ ) (Table 4). We also wanted to assess with binary logistic regression model whether intraoperative SAS was better predictor for the 30-days major complications than the following patients' predictor variables ASA physical status, age and gender (Table 5). Binary logistic regression model was statistically significant ( $\chi^2$  test=9,82, df=4,  $P=0,043$ ) and explained 33,1% of dependent variable variance. SAS score  $\leq$ 4 was only predictor variable that significantly influence on belonging in major 30-days postoperative complications group: OR=14,24 (95% CI: 1,84-110,32) controlled for all other variables in the model (Table 5). Moreover, the length of operative delay was significantly associated with the 6-months major complications when Mann-Whitney test was applied ( $p=0.015$ ). Those 4 patients who developed 6-months major complications had median length of operative delay for 7 days (IQR 6.25-7). It is relevant to emphasize that these patients did not develop 30-days major complications.

## Discussion

So far many studies have observed value of this 10-point intra- and postoperative surgical outcome score in many surgical subspecialties. Although this score has been validated in selected orthopaedic procedures (10,11,12,13,14) we also wanted to observe the utility of the SAS in the trauma hip fracture surgery patients. Hip fracture surgery repairs are operations of different magnitude. Together with poor preoperative medical condition of these patients, many comorbidities, and reduced cardiopulmonary reserves, preoperative variables like age and ASA physical status sometimes are not enough to prognose early and late postoperative patient's course and the need for the ICU surveillance. Our study demonstrated that the SAS  $\leq$ 4 was the only significant predictor for the 30-days major complications development when compared with the patient's gender, age and ASA physical status. The cases with the SAS $\leq$ 4 had 14.4 times greater odds than the cases with the SAS $\geq$ 5 for developing 30-days major complications. Our result is almost consistent with other studies in which these poor-scoring patients were 16 times more likely to develop major

complications (5,8). This is very important because the median postoperative hospital stay in our study was 10 days. In addition, our research also observed that the operative delay was associated with the 6-months major complications development. Cases who developed 6-months major complications had operative delay approximately for 7 days. Reasons for such a long operative delay were the preoperative poor medical patient's condition and time to optimize it, additional diagnostic test requirements, availability of the ICU postoperative surveillance. However, every clinician should be aware that operative delay of more than 48 hours, especially for low-risk and younger patients, is associated with increased short-term (30-days) and mid term (1-year) mortality, as well as with prolonged hospital stay of these patients (15,16,17). Yet the mortality rate for this fragile geriatric population approaches expected mortality approximately 6-months postoperatively according to the research (18). So the hip fracture patients with the  $SAS \leq 4$  and the operative delay of more than 48 hours every clinician should recognise as red alarm warning for not only the postoperative intensive hospital care but also the postoperative 6-months surveillance to prevent worse outcome.

This surgical score is also a reliable intraoperative patient guiding tool for all the members of the surgical team. There was no difference in the MAP values between the  $SAS \leq 4$  and  $SAS \geq 5$  groups probably due to applied vasoactive bolus doses support intraoperatively to avoid current hypotension. This refers to a good intraoperative blood pressure control. But the mentioned SAS groups significantly varied by the intraoperative HR and blood loss. In our study the lower HR and estimated blood loss were associated with higher SAS values and the better primary patient's outcome. The SAS value would be higher by avoiding higher HR, hypotension and applying surgical technique to minimize the bleeding. Hence, we also observed that maintaining intraoperative stability of the vital signs and controlling the bleeding are relevant predictors of the patient's final outcome (19,20).

Our study has several limitations. First, the SAS was tested only in the small size trauma hip fracture patients while its importance and implementation in the postoperative surgical outcome risk grading in other trauma patients is still not recognized. In addition retrospective research can have lots of falsity and underreporting data. Second, the SAS variables were gained from the handwritten anaesthesia records which are not reliable as electronic versions. Also, blood loss estimation could be questionable. Yet, the studies have shown that the SAS blood loss estimation categories match the observer's blood loss volume accurately, and obtain more reliability if estimation is made by the anaesthesiologist (21,22). Therefore, in comparison with the obstetric Apgar score the SAS variables are nothing less reliable than the obstetric Apgar score parameters (23). Moreover, another limitation of our research was that all-cause mortality was the primary endpoint, without specific secondary endpoints such as cardiovascular, neurologic, or infectious events. That is



because recent studies have shown that this secondary endpoints can not accurately predict mortality therefore are not recommended for the studies which aim is to decrease mortality (24,25,26). Another weakness of the score is that it can not be used to compare physicians or institutions because it greatly depends on the patient's preoperative condition and yet not established in our country risk-adjustment National Surgical Quality Improvement Program model. In spite of the limitations, we think that a future prospective research about the SAS utility in the all trauma procedures on a large sample size is necessary.

## **Conclusion**

To conclude, intraoperative patient's course has an impact on the postoperative outcome. Thereby implementation of this simple surgical outcome score in everyday routine practise for the trauma hip fracture patients gives us objective, immediate, feed-back information about patient's clinical condition at the end of the procedure. However, our study showed that every hip fracture patient with the  $SAS \leq 4$  should postoperatively have ICU surveillance. Likewise enables us to stratify patients with the higher than average odds for 30-days major complications. It is also a tool which facilitates good interdisciplinary operative and postoperative communication between the team members, and routes us to apply suitable and quality postoperative care management for the individual patient in every aspect.

## References

1. Kang JW, Kim KJ, Lee SK, Kim J, Jeung SW, Choi HG. Predictors of mortality in patients with hip fractures for persons aging more than 65 years old. *International Journal of Bio-Science and Bio-Technology* 2013;5:27-34.
2. Donegan DJ, Gay AN, Baldwin K, Morales EE, Esterhai Jr JL, Mehta S. Use of medical comorbidities to predict complications after hip fracture surgery in the elderly. *The Journal of Bone & Joint Surgery* 2010; 92:807–813.
3. Egol KA, Strauss EJ. Perioperative considerations in geriatric patients with hip fracture: what is the evidence? *J Orthop Trauma* 2009;23:386-94. doi: 10.1097/BOT.0b013e3181761502.
4. Reynolds PQ, Sanders NW, Schildcrout JS, Mercaldo ND, St Jacques PJ. Expansion of the surgical Apgar score across all surgical subspecialties as a means to predict postoperative mortality. *Anesthesiology* 2011;114:1305-12. doi: 10.1097/ALN.0b013e318219d734.
5. Gawande AA, Kwaan MR, Regenbogen SE, Lipsitz SA, Zinner MJ. An Apgar score for surgery. *J Am Coll Surg* 2007;204:201-8.
6. Reynolds PQ, Sanders NW, Schildcrout JS, Mercaldo ND, Jacques PJ. Expansion of the Surgical Apgar Score across all surgical subspecialties as a means to predict postoperative mortality. *Anesthesiology* 2011;114:1305-12.
7. ASA Physical Classification System, American Society of Anesthesiologists, <http://www.asahq.org/clinical/physicalstatus>.
8. Regenbogen SE, Ehrenfeld JM, Lipsitz SR, Greenberg CC, Hutter MM, Gawande AA. Utility of the Surgical Apgar Score. Validation in 4119 patients. *Arch Surg* 2009;144:30-36.
9. Ziewacz JE, Davis MC, Lau D, El-Sayed AM, Regenbogen SE, Sullivan SE et al. Validation of the surgical Apgar score in a neurosurgical patient population. *J Neurosurg* 2013;118:270-9. doi: 10.3171/2012.10.JNS12436. Epub 2012 Nov 2.
10. Haynes AB, Regenbogen SE, Weiser TG, Lipsitz SR, Dziekan G, Berry WR, et al: Surgical outcome measurement for a global patient population: validation of the Surgical Apgar Score in 8 countries. *Surgery* 2011;149:519–524.
11. Prasad SM, Ferreria M, Berry AM, Lipsitz SR, Richie JP, Gawande AA, et al: Surgical Apgar outcome score: perioperative risk assessment for radical cystectomy. *J Urol* 2009;181:1046–1053.

12. Regenbogen SE, Ehrenfeld JM, Lipsitz SR, Greenberg CC, Hutter MM, Gawande AA. Utility of the surgical apgar score: validation in 4119 patients. *Arch Surg* 2009;144:30–37.
13. Wuerz TH, Regenbogen SE, Ehrenfeld JM, Malchau H, Rubash HE, Gawande AA, et al: The Surgical Apgar Score in hip and knee arthroplasty. *Clin Orthop Relat Res* 2011;469:1119–1126.
14. Zigelboim I, Kizer N, Taylor NP, Case AS, Gao F, Thaker PH, et al: “Surgical Apgar Score” predicts postoperative complications after cytoreduction for advanced ovarian cancer. *Gynecol Oncol* 2010;116:370–373.
15. Zuckerman JD, Skovron ML, Koval KJ, et al. Postoperative complications and mortality associated with operative delay in older patients who have a fracture of the hip. *J Bone Joint Surg Am.* 1995;77:1551–1556.
16. Siegmeth AW, Gurusamy K, Parker MJ. Delay to surgery prolongs hospital stay in patients with fractures of the proximal femur. *J Bone Joint Surg Br.* 2005;87:1123–1126.
17. Shiga T, Wajima Z, Ohe Y. Is operative delay associated with increased mortality of hip fracture patients? Systematic review, meta-analysis, and meta-regression. *Can J Anaesth.* 2008;55:146-54. doi: 10.1007/BF03016088.
18. Magaziner J, Simonsick EM, Kashner TM, Hebel JR, Kenzora JE. Survival experience of aged hip fracture patients. *Am J Public Health.* 1989;79:274-8.
19. Hartmann B, Junger A, Röhrig R, et al. Intra-operative tachycardia and perioperative outcome. *Langenbecks Arch Surg* 2003;388:255–260.
20. Pasternak LR. Risk assessment in ambulatory surgery: challenges and new trends. *Can J Anesthesia* 2004;51:R1–R5.
21. Delilkan AE. Comparison of subjective estimates by surgeons and anaesthetists of operative blood loss. *Br Med J* 1972;2:619–621.
22. Gardiner AJS, Dudley HAF. The measurement of blood loss at operation. *Br J Anesth* 1962;34:653–656.
23. Finster M, Wood M. The Apgar score has survived the test of time. *Anesthesiology* 2005;102:855-857.
24. Cannon CP: Clinical perspectives on the use of composite endpoints. *Control Clin Trials* 1997;18:517-29; discussion 546-9.
25. O’Neil RT: Secondary endpoints cannot be validly analyzed if the primary endpoint does not demonstrate clear statistical significance. *Control Clin Trials* 1997;18:550-6; discussion 561-7.

26. POISE Study Group, Devereaux PJ, Yang H, Yusuf S, Guyatt G, Leslie K, Villar JC, Xavier D, Chrolavicius S, Greenspan L, Pogue J, Pais P, Liu L, Xu S, Malaga G, Avezum A, Chan M, Montori VM, Jacka M, Choi P: Effects of extended-release metoprolol succinate in patients undergoing non-cardiac surgery ( POSE trial): A randomised controlled trial. *Lancet* 2008;371:1839-47.

**Abbreviations:** American Society of Anaesthesiologists (ASA); Surgical Apgar Score (SAS); Estimated blood loss (EBL); Heart rate (HR); Mean arterial pressure (MAP)

**Acknowledgements:** The authors certify that there is no known financial or non-financial interest associated with this publication. The authors declare that they have no competing interest.

**Figure 1. Calculating the 10-point Surgical Apgar Score<sup>1</sup>**

<b>Surgical Apgar Score Points</b>					
	<b>0 points</b>	<b>1 point</b>	<b>2 point</b>	<b>3 points</b>	<b>4 points</b>
<b>Estimated blood loss (ml)</b>	<b>&gt; 1000</b>	<b>601-1000</b>	<b>101-600</b>	<b>≤100</b>	<b>—</b>
<b>Lowest mean arterial pressure (mmHg)</b>	<b>&lt; 40</b>	<b>40-54</b>	<b>55-69</b>	<b>≥ 70</b>	<b>—</b>
<b>Lowest heart rate (beats/min)</b>	<b>&gt; 85</b>	<b>76-85</b>	<b>66-75</b>	<b>56-65</b>	<b>≤ 55*</b>

<sup>1</sup>The Surgical Apgar score is computed at the end of the operation from the anaesthesia records as the sum of the 3 intraoperative variables: estimated blood loss, lowest mean arterial pressure, and lowest heart rate.

\*Occurrence of pathologic bradyarrhythmia including sinus arrest, atrioventricular block or dissociation, junctional or ventricular escape rhythms, and asystole, also receives 0 points for lowest heart rate.

**Table 1. Baseline perioperative characteristics of hip fracture patients**


		N	%
Gender	Male	8	18,6%
	Female	35	81,4%
ASA class	2	3	7,0%
	3	35	81,4%
	4	5	11,6%
Pulmonary disease <sup>1</sup>	No	28	65,1%
	Yes	15	34,9%
Cardiovascular disease <sup>2</sup>	No	13	30,2%
	Yes	30	69,8%
Renal disease <sup>3</sup>	Ne	37	86,0%
	Da	6	14,0%
Coagulation disorders <sup>4</sup>	No	35	81,4%
	Yes	8	18,6%
Estimated blood loss <sup>l</sup> ml	>1000 ml	8	18,6%
	601-1000 ml	9	20,9%
	101-600 ml	17	39,5%
	<=100 ml	9	20,9%
Type of operation	Dynamic hip screw	18	41,9%
	Hemiarthroplasty	19	44,2%
	Total hip replacement	1	2,3%
	Osteosynthesis sec AO	5	11,6%
Operative diagnosis	Pertrochanteric fracture	19	44,2%
	Femoral neck fracture	20	46,5%
	Subtrochanteric fracture	4	9,3%
Anaesthesia technique	General	27	62,8%
	Spinal	16	37,2%
Surgery urgency	Elective	0	,0%
	Expedited	40	93,0%
	Urgent	3	7,0%
Vasoactive support	No	37	86,0%
	Yes	6	14,0%
Major complications, 30-days	No	35	81,4%
	Yes	8	18,6%
Hospital mortality	No	42	97,7%
	Yes	1	2,3%
30-days mortality	No	38	88,4%
	Yes	5	11,6%
Major complications, 6-months	No	39	90,7%
	Yes	4	9,3%

<sup>1</sup> Pulmonary comorbidity was defined as pneumonia, mechanical ventilator dependency, and preexisting chronic obstructive pulmonary disease. <sup>2</sup>Cardiovascular comorbidity included earlier myocardial infarction, angina pectoris, congestive heart disease, coronary revascularization, peripheral vascular disease and anamnesis of stroke and transient ischaemic attack. <sup>3</sup> Renal comorbidity included history of acute or chronic renal disease. <sup>4</sup>Coagulation comorbidity comprised hereditary and acquired coagulation disorder, as well as the use of anticoagulation agents such as warfarin, acetylsalicylic acid and clopidogrel.

**Table 2. Perioperative hip fracture patients' characteristics expressed as mean  $\pm$  SD and median (25%-75% IQR) after applying t-test.**

	Mean $\pm$ SD	Percentiles	p value*
		Median (IQR)	
Age, years	80.30 $\pm$ 10.09	82 (75-87)	0.498
ASA class	3.05 $\pm$ 0.43	3 (3-3)	
Preoperative haemoglobin, g/L	121.51 $\pm$ 19.61	122 (111-132)	0.203
Preoperative platelet count, x 10 <sup>9</sup> L	235.79 $\pm$ 94.19	221 (182-265)	0.607
Preoperative prothrombin time	0.95 $\pm$ 0.24	1.02 (0.9-1.11)	0.533
Operative delay, days	3.84 $\pm$ 2.51	3 (2-6)	0.336
Lowest MAP, mmHG	65.13 $\pm$ 15.85	66 (53-76)	0.101
Lowest HR, beats/min	72.56 $\pm$ 16.49	70 (65-85)	0.007
SAS value	5.53 $\pm$ 1.79	6 (4-7)	
Operative duration, min	104.76 $\pm$ 61.19	90 (70-120)	0.055
Postoperative haemoglobin, g7L	109.23 $\pm$ 16.72	109 (98-123)	0.151
ICU length of stay, days	4 $\pm$ 5.66	2 (2-3)	0.315
Hospital length of stay, days	11.74 $\pm$ 9.27	10 (8-13)	0.495

\*The p values were obtained after applying Mann-Whitney test to compare SAS value with perioperative variables. P value <0.05 was considered significant.



**Figure 2. Receiver operating characteristics curve for the SAS value  $\leq 4$  and 30-days major complications. The SAS value  $\leq 4$  was a significant predictor for the 30-days major surgical complications with 80% specificity. ROC AUC 0.743, SE 0.0957, 95% CI: 0.587-0.864,  $p=0.0111$  for Area =0.5.**



**Table 3. After the SAS cut-off value  $\leq 4$  patients were divided in two SAS groups and their perioperative variables were compared applying Pearson Chi-Square Test.\* P value  $<0.05$  was considered significant.**

		SAS: groups				p value
		$\leq 4$		$> 4$		
		N	%	N	%	
Gender	Male	3	25.0%	5	16.1%	0.503
	Female	9	75.0%	26	83.9%	
ASA class	2	0	0.0%	3	9.7%	0.465
	3	11	91.7%	24	77.4%	
	4	1	8.3%	4	12.9%	
Pulmonary disease	No	9	75,0%	19	61.3%	0.398
	Yes	3	25,0%	12	38.7%	
Cardiovascular disease	No	4	33.3%	9	29,0%	0.783
	Yes	8	66.7%	22	71,0%	
Renal disease	No	9	75,0%	28	90.3%	0.193
	Yes	3	25,0%	3	9.7%	
Coagulation disorders	No	11	91.7%	24	77.4%	0.282
	Yes	1	8.3%	7	22.6%	
Estimated blood loss, ml	$>1000$ ml	5	41.7%	3	9.7%	0.016
	601-1000 ml	4	33.3%	5	16.1%	
	101-600 ml	3	25,0%	14	45.2%	
	$\leq 100$ ml	0	0,0%	9	29,0%	
Type of operation	Dynamic hip screw	3	25,0%	15	48.4%	0.451
	Hemiarthroplasty	7	58.3%	12	38.7%	
	Total hip replacement	0	0,0%	1	3.2%	
	Osteosynthesis sec AO	2	16.7%	3	9.7%	
Operative diagnosis	Pertrochanteric fracture	3	25,0%	16	51.6%	0.054
	Femoral neck fracture	6	50,0%	14	45.2%	
	Subtrochanteric fracture	3	25,0%	1	3.2%	
Anaesthesia technique	General	8	66.7%	19	61.3%	0.744
	Spinal	4	33.3%	12	38.7%	
Operative urgency	Elective	0	0,0%	0	0,0%	0.121
	Expedited	10	83.3%	30	96.8%	
	Urgent	2	16.7%	1	3.2%	
Vasoactive support	No	8	66.7%	29	93.5%	0.022
	Yes	4	33.3%	2	6.5%	
Major complications, 30-days	No	7	58.3%	28	90.3%	0.016
	Yes	5	41.7%	3	9.7%	
Hospital mortality	No	11	91.7%	31	100,0%	0.104
	Yes	1	8.3%	0	0,0%	
30-days mortality	Ne	10	83.3%	28	90.3%	0.521
	Da	2	16.7%	3	9.7%	
Major complications, 6-months	No	10	83.3%	29	93.5%	0.301
	Yes	2	16.7%	2	6.5%	

**Table 4. Correlation of the SAS and perioperative patients variables using Spearman Correlation Coefficient. P (2-tailed) value <0.05 was considered significant.**

<b>Variables</b>	<b>rho*</b>	<b>p value (2-tailed)</b>
Age, years	-0.002	0.992
ASA class	-0.037	0.814
Preoperative haemoglobin, g/L	0.111	0.477
Preoperative platelet count, x 10 <sup>9</sup> L	-0.051	0.746
Preoperative prothrombin time	0.171	0.274
Operative delay, days	-0.076	0.627
Estimated blood loss, ml	0.600	<0.001
Lowest mean arterial pressure, mmHG	0.256	0.098
Lowest heart rate, beats/min	-0.509	<0.001
Postoperative haemoglobin, g/L	0.230	0.138
Operative duration, min	-0.375	0.014
ICU length of stay, days	0.030	0.849
Hospital length of stay, days	-0.057	0.719

\*rho is Correlation Coefficient. – sign mens negative correlation, no sign means positive correlation.

**Table 5. Binary logistic regression model which included 4 predictor variables ( ASA score, SAS score,age and gender) has been made to assess prediction to group with 30 days postop major complications as a binary dependent variable (0=without major complications and 1=with complicationa). Regression model was statistically significant (X2 test=9,82, df=4, P=0,043) and explained 33,1% of dependent variable variance. SAS score <=4 was only predictor variable that significantly influence on belonging in major compicatin group: OR=14,24 (95% CI: 1,84-110,32) controlled for all other variables in the model.**

	B	S.E.	Wald	df	OR	95% CI		P
						Lower	Upper	
ASA score	-1,50	1,24	1,47	1	0,22	0,02	2,52	0,226
SAS score <=4	2,66	1,04	6,47	1	14,24	1,84	110,32	0,011
Age (years)	-0,05	0,05	1,00	1	0,95	0,86	1,05	0,317
Female gender	1,37	1,35	1,03	1	3,93	0,28	55,29	0,311