



Središnja medicinska knjižnica

Wolstencroft, E. C., Šimić, G., thi Man, N., Holt, I., Lam, Le T., Buckland, P. R., Morris, G. E. (2007) *Endosomal location of dopamine receptors in neuronal cell cytoplasm*. *Journal of Molecular Histology*, 38 (4). pp. 333-340.

The original publication is available at www.springerlink.com

<http://www.springerlink.com/content/353t21168pq07477/>

<http://dx.doi.org/10.1007/s10735-007-9106-5>

<http://medlib.mef.hr/323>

University of Zagreb Medical School Repository

<http://medlib.mef.hr/>

Elizabeth C. Wolstencroft¹, Goran Šimić², Nguyen thi Man^{1,3}, Ian Holt^{1,3}, Le Thanh Lam^{1,3}, Paul R. Buckland⁴, Glenn E. Morris^{1,3}

Endosomal location of dopamine receptors in neuronal cell cytoplasm

¹Wolfson Centre for Inherited Neuromuscular Disease, RJA Orthopaedic Hospital, LMARC Building, Oswestry SY10 7AG, UK

²Department of Neuroscience, Croatian Institute for Brain Research, University Medical School, Salata 12, Zagreb, Croatia

³Institute of Science and Technology in Medicine, Keele University, Keele ST4 7QB, UK

⁴Department of Psychological Medicine, University of Wales College of Medicine, Heath Park, Cardiff, UK

Correspondence to:

Glenn E. Morris, Wolfson Centre for Inherited Neuromuscular Disease, RJA Orthopaedic Hospital, LMARC Building, Oswestry SY10 7AG, UK, e-mail: glenn.morris@rjah.nhs.uk

Abstract

Five subtypes of dopamine receptor exist in two subfamilies: two D1-like (D1 and D5) and three D2-like (D2, D3 and D4). We produced novel monoclonal antibodies against all three D2-like receptors and used them to localize receptors in Ntera-2 (NT-2) cells, the human neuronal precursor cell line. Most of the immunostaining for all three receptors colocalized with mannose-6-phosphate receptor, a marker for late endosomes formed by internalization of the plasma membrane. This result was obtained with antibodies against three different epitopes on the D3 receptor, to rule out the possibility of cross-reaction with another protein, and controls without primary antibody or in the presence of competitor antigen were completely negative. In rat cerebral cortex and hippocampus, some of the dopamine receptor staining was found in similar structures in neuronal cell cytoplasm. Only some of the neurons were positive for dopamine receptors and the pattern was consistent with previously-reported patterns of innervation by dopamine-producing neurons. Endosomal dopamine receptors may provide a useful method for identifying cell bodies of dopamine-responsive neurons to complement methods that detect only active receptors in the neuronal cell membrane.

Keywords

G protein-coupled receptor, endosome, NT-2 cells, Ntera-2, monoclonal antibody, brain, hippocampus, cerebellum, cerebral cortex, mannose-6-phosphate receptor

Introduction

Dopamine receptors are members of the G protein-coupled receptor (GPCR) family, which initiate their biological function by coupling to GTP regulatory proteins (G proteins). In humans, five dopamine receptor subtypes, DRD1 (Monsma et al.

1990), DRD2 (Bunzow et al. 1988), DRD3 (Sokoloff et al. 1990), DRD4 (van Tol et al. 1991) and DRD5 (Sunahara et al. 1991) have been identified. The D1-like receptors, DRD1 and DRD5, stimulate adenylyl cyclase (Monsma et al. 1990; Albert et al. 1990) whereas the D2-like receptors, DRD2, DRD3 and DRD4, are coupled to inhibitory G proteins and their stimulation leads to inhibition of adenylyl cyclase (Sokoloff et al. 1990). All five have an extracellular amino terminus, seven transmembrane domains and an intracellular carboxyl terminus. Abnormalities in dopamine neurotransmission are implicated in the aetiology of neurological and psychiatric disorders including schizophrenia (Williams et al. 1998) and Parkinson's disease (Oliveri et al. 2000).

The distribution of D2-like dopamine receptor mRNAs in brain has been studied by *in situ* hybridization (Sokoloff et al. 1990; Mengod et al. 1989) and radioligand binding studies (Martres et al. 1985; Bouthenet et al. 1987; Camps et al. 1989) have revealed the distribution of biologically active receptors, including the use of subtype selective ligands (Levesque et al. 1992; Herroelen et al. 1994; Hall et al. 1996; Bancroft et al. 1998). Most immunohistochemical studies of D2-like receptors so far have used only a single antibody, with the notable exception of two studies of DRD4 that used antibodies against two epitopes (Defagot et al. 1997; Mauger et al. 1998). Use of antibodies against two or more epitopes is the best safeguard against cross-reaction, especially with monoclonal antibodies (Morris 1998). The reported sizes of D2-like dopamine receptors on western blots have varied widely in the literature (36–95 kDa) and cross-reaction with other proteins has not always been ruled out.

In the present study, we have used a new panel of antibodies against three different DRD3 epitopes to determine the cellular and subcellular localization of DRD3 in rat brain and Ntera-2 cells. Additional monoclonal antibodies against specific peptide sequences in DRD2 and DRD4 were used to support and confirm the subcellular localization of D2-like dopamine receptors. We suggest that the presence of D2-like dopamine receptors in cytoplasmic vesicles could provide a useful means to identify the cell bodies of dopamine-responsive neurons, since neuronal extensions with dopamine receptors often project far into other regions of the brain.

Methods

Antibody production

For monoclonal antibodies, Balb/c mice were immunized with peptides conjugated to bovine serum albumin (BSA) through cysteine residues using MBS (3-maleimidobenzoic acid N-succinimidyl ester). Suitable human peptide sequences were identified on the basis of specificity for each subtype and antigenicity. The peptides were synthesized by Alta Bioscience (University of Birmingham, UK). Hybridoma fusions with the Sp2/0 mouse myeloma cell line were performed as described previously (Nguyen and Morris 1996). Hybridoma culture supernatants were screened by enzyme linked immunosorbent assay (ELISA) using plates coated with the respective unconjugated peptide. ELISA positive wells were further screened by Western blotting on total Ntera-2 protein extract and total brain (rat and human) extract.

A rabbit polyclonal antiserum raised against the same N-terminal DRD3 peptide was obtained from Calbiochem. A polyclonal antiserum was produced against a DRD3 Cterminal recombinant protein (amino acids 205–400) by immunization of Balb/c mice. The corresponding cDNA was obtained from a full-length human DRD3 construct by PCR and was cloned into pET 32a expression vector (Novagen, Darmstadt, Germany). The expressed protein was purified for immunization by affinity chromatography using His.Bind resin (Novagen, Darmstadt, Germany) and used to immunize Balb/c mice. All animal procedures were performed under UK license.

Enzyme linked immunosorbent assay

A 96 well PVC microtitre plate (Dynatech, Billingshurst, UK) was coated with 50 μ l per well of the peptide at a concentration of 5–10 μ g/ml in PBS, and incubated at room temperature for 1 h. The antigen was then removed and the plate was washed 3 times with incubation buffer (IB, PBS + 1% Triton X-100). 300 μ l of a 4% solution of skimmed milk powder in IB was added to each well for 30 min at room temperature. After washing 3 times with IB, 50 μ l of culture supernatant was added at room temperature for 1 h. The primary antibody was removed and the plate washed 4 times with IB. Horseradish peroxidaseconjugated rabbit anti-mouse Ig (DAKO patts, Denmark; 50 μ l of 1/1,000 in dilution buffer: 0.1% BSA, 1% horse serum and 1% fetal calf serum in IB) was then added and incubated at room temperature for 1 h. The plate was washed 4 times with IB. One hundred and fifty microlitres of substrate solution (0.4 mg/ml o-phenylene-diamine in 25 mM phosphate-citrate buffer pH 5.0 with 0.1% H₂O₂) was added until the colour developed. The reaction was then stopped with 50 μ l of 3M H₂SO₄. Samples were read using a 492 nm filter on a Titertek Multiscan Plate reader.

Western blotting

Human brain samples were provided by the Medical Research Council Brain Bank, Kings College, London, UK. Total protein extracts of human or rat brain were made by Dounce homogenization in four volumes of SDS extraction buffer (10% SDS, 10% EDTA, 5% 2-mercaptoethanol, 10% glycerol, 50mM Tris-HCl, pH 7.0) and clarified by centrifugation at 160,000g for 30 min. After SDS-PAGE on 10% polyacrylamide gels, proteins were transferred to nitrocellulose electrophoretically overnight. After blocking for 1 h in 3% skimmed milk powder in incubation buffer (0.05% Triton X-100 in PBS), blots were incubated with mAb (1/100 dilution of culture supernatant in incubation buffer/1% horse serum/1% fetal calf serum/0.1% BSA) for 1 h. After washing 3 times in PBS, blots were incubated with biotinylated anti-mouse Ig and a peroxidase-avidin detection reagent (Vectastain ABC Elite, Vector Laboratories, Peterborough, UK) according to the manufacturer's instructions. Blots were developed with diaminobenzidine in 25mM phosphatecitrate buffer (pH 5.0), 0.012% H₂O₂.

Cell culture

The human embryonal carcinoma cell line, Ntera-2 (Andrews et al. 1984), also known as NT-2, was cultured in Dulbecco's MEM with Glutamax-I (Invitrogen,

Paisley, UK) and 10% fetal bovine serum (Invitrogen, UK). Cells were seeded onto coverslips at $5 \cdot 10^5$ cells and, after culture for 48 h, were fixed in 50:50 acetone–methanol for 5 min.

Preparation of rat brain sections

A whole brain was removed from an adult Sprague-Dawley rat immediately post-mortem and rinsed in ice-cold isotonic saline for 1 h. It was then mounted and frozen in isopentane cooled in liquid nitrogen and sectioned dorsolaterally into 5 μ m frozen sections. The sections were mounted on microscope slides in the same orientation and allowed to air dry.

Immunofluorescence microscopy

Cells and tissue sections were fixed using 50% acetone/50% methanol. The primary antibody was diluted in PBS/1% horse serum/1% fetal calf serum and incubated on the slide for 1 h at room temperature. After washing with PBS, FITC-conjugated rabbit anti-(mouse Ig) (1/50: Vector Labs, Burlingame, USA) in PBS/1% horse serum/1% fetal calf serum was applied for 30 min. After counterstaining nuclei with ethidium bromide, slides were mounted in 10% glycerol in PBS and photographed using a Leica DMRB epifluorescence photomicroscope with 10 \times or 63 \times objectives. For double labelling, fixed cells were incubated sequentially with rabbit antisera and mouse mAb for 1 h, followed by a mixture of FITC-conjugated anti-(mouse Ig) (1/50 dilution) (Vector Labs, Burlingame, CA, USA) and TRITC-conjugated anti-(rabbit Ig) (1/100 dilution) (DAKO, Glostrup, Denmark) for 30 min. Confocal images were captured using a Zeiss LSM510 Meta confocal system.

Results

Production of antibodies that recognize 50 kDa dopamine receptors in human brain

A panel of monoclonal antibodies (Table 1) was raised against the three D2-like dopamine receptors using BSAconjugated, subtype-specific peptides corresponding to the human DRD3 N-terminal domain (ASLSQLSSH-C; two mAbs MANDOP31 and 32), the DRD3 third cytoplasmic domain (RIYVVLKQR-C; three mAbs MANDOP33–35), the DRD2 third cytoplasmic domain AARRAQELEMEC; two mAbs MANDOP21 and 22) and the DRD4 third extracellular domain (DVRQRDPAVC; one mAb MANDOP41). Hybridoma fusions were screened by ELISA with unconjugated peptide and by Western blotting using total human and rat brain extracts. Figure 1a shows three mAbs against both DRD3 epitopes (MANDOP31, 32 and 35) that recognize a band of the expected size (50 kDa) in a human brain extract. The two mAbs against DRD2 and one against DRD4 also recognized a protein of the same size (Fig. 1b).

The DRD2 and N-terminal DRD3 peptide immunogens show no sequence similarity to the other DRD subtype sequences, but the DRD4 peptide does have four of ten amino acids identical to DRD3 only, so the possibility of cross-reaction with DRD3 cannot be entirely ruled out. A C-terminal DRD3 fragment (amino acids 205–400) expressed from a pET32 plasmid and purified by affinity chromatography was used to produce a polyclonal antiserum that will recognize a

third DRD3 epitope for confirmation of specificity. Subcellular fractionation was carried out on total Ntera-2 cell extract by a series of high-speed centrifugations. The 50 kDa protein recognized by MANDOP35 was absent from cytosolic fractions until extracted by Triton X-100, consistent with membrane-associated DRD3 (Fig. 1c).

Most of the D2-like dopamine receptors in cultured Ntera-2 cells are found in endosomal vesicles in the cytoplasm

Ntera-2 cells exhibit properties of a committed neuronal precursor at an early stage of development (Andrews et al. 1984) and have been shown by reverse transcriptase-polymerase chain reaction and functional studies to express DRD2 and DRD3 dopamine receptors (Hurlbert et al. 1999; Sodja et al. 2002). Using immunofluorescence microscopy, all dopamine receptors were found almost exclusively in large cytoplasmic vesicles in Ntera-2 cells (Fig. 2a). No such staining was seen in HeLa cells or human skin fibroblasts. To show that this staining pattern was not due to a cross-reaction, double label immunofluorescence microscopy was carried out using a mouse monoclonal antibody against the DRD3 third cytoplasmic domain (MANDOP34), and a commercial rabbit polyclonal antiserum against the DRD3 N-terminus and their co-localization was complete (Fig. 2b). All three monoclonal DRD3 epitopes gave the same result and there was no staining in the absence of primary antibody (inset: Fig. 2a). Antibody staining of vesicles was also prevented by the peptide antigen (see Fig. 3). A double label using the same commercial rabbit antiserum against DRD3 and a commercial mouse monoclonal antibody against mannose-6-phosphate receptor, a marker for late endosomes, showed co-localization (Fig. 2c), suggesting that the cytoplasmic vesicles are a late stage in the pathway for receptor internalization and turnover via lysosomes.

Some dopamine receptors were detected at the expected plasma membrane sites

One would normally expect to find D2-like dopamine receptors in the plasma membrane, close to synapses with dopamine-producing neurons, and Fig. 2d shows a section of rat basal ganglia with DRD3 staining of dendrite-like processes. Similar staining was observed in other areas of the brain, including cortical layer V of cerebral cortex, where dopaminergic innervation is also expected (not shown). Identification and proper characterization of different dendritic processes is a complex neuroscience study beyond the scope of this paper, but we show Fig. 2d to emphasize that our mAbs may detect dopamine receptors in their functional location in plasma membranes, as well as in endosomes. Indeed, DRD3 receptors are sometimes detectable in the plasma membrane of Ntera-2 cells (Fig. 2e), although cell culture appears to favour location in endosomes.

Endosome-like staining by dopamine receptor antibodies can be used to identify specific subsets of rat brain neurons

Figure 3a shows that all three receptors are also present as cytoplasmic vesicles in neurons of cortical layers V and VI of rat cerebral cortex. Antibody staining of the vesicles was prevented by pre-incubation with peptide antigen (see inset for DRD3 control). The DRD2 mAb shows additional staining in layer I, but, with only one mAb available, we cannot rule out cross-reaction with a non-DRD protein. Cross-

reaction of MANDOP41 with DRD3 cannot be entirely ruled out. Figure 3b shows a section through the hippocampus and dentate gyrus. Cytoplasmic vesicles containing D2-like dopamine receptors were found in CA3 and other neurons, but were absent from CA1 neurons and from granule cells of the dentate gyrus. Similar endosomelike vesicles were observed in cerebellum (Purkinje neurons) and brain stem (large bipolar neurons) (data not shown). This shows that DRD-positive vesicle staining could be a useful means of identifying the cell bodies of neurons that are making different dopamine receptors.

Discussion

Immunolocalization studies using monoclonal antibodies against the D2-like dopamine receptors have shown an endosomal distribution in the Ntera-2 cell line. This distribution pattern is not a staining artifact, since it is completely dependent on the presence of the primary DRD-specific mAb and is blocked by pre-incubation with antigen. We are confident of this localization for DRD3 since an identical pattern was produced by mAbs against two different epitopes in the N-terminal and third cytoplasmic domains, as well as by an antiserum against the C-terminal region and a commercial rabbit antiserum. This confidence was reinforced (a) by the fact that mAbs against DRD2 (different part of third cytoplasmic domain) and DRD4 (third extracellular domain) stained these vesicles, suggesting that all three D2-like dopamine receptors are co-distributed in these vesicles in Ntera-2 cells, and (b) by the observation of similar vesicles in rat brain sections. Because unique peptide sequences were chosen as immunogens for each subtype, cross-reaction between D2-like dopamine receptors is hardly possible, except for DRD4 where the possibility of cross-reaction with DRD3 cannot be entirely ruled out. Cross-reaction with some functionally-unrelated protein can be ruled out when antibodies against two or more epitopes are used (i.e. DRD3) but not when only one mAb is available (e.g. DRD2 in Fig. 3). According to current ideas about GPCR trafficking in general (Dohlman et al. 1991), and that of dopamine receptors in particular (Ng et al. 1994), receptors are internalized in “early endosomes” from which they may be recycled to the plasma membrane or targeted for turnover in “late endosomes”. It is interesting to note that the dopamine transporter protein, which regulates dopamine receptor activation, is known to be internalized into endosomes by a clathrin-dependent mechanism (Sorkina et al. 2005). Internalization and recycling of GPCRs involves arrestin and a phosphorylation/dephosphorylation cycle (Ng et al. 1994; Ferguson et al. 1996). A number of transfection studies of DRD1 and DRD2 have shown a predominantly cytoplasmic localization for DRD2, while DRD1 is localized in the plasma membranes after transfection (Ng et al. 1994; Vickery and von Zastrow 1999; Prou et al. 2001). Agonist stimulation, however, also results in internalization of DRD1 (Ng et al. 1994; Vickery and von Zastrow 1999). Internalization of the receptor following ligand binding is known to occur for a number of GPCRs, including the D2 type receptors, and this leads to a lowering of surface receptor density (Ferguson 2001; Sun et al. 2003). Agonist-independent internalization of D2 type receptors may be occurring in Ntera-2 cells under the culture conditions used in this study.

Previous immunohistological studies of dopamine receptors in the brain have shown the presence in neuronal cells of DRD1 and 2 (Levey et al. 1993) and DRD4 (Defagot et al. 1997; Wedzony et al. 2000; Rivera et al. 2002) but subcellular

distribution of the receptors was not described. Our present antibody studies of rat brain sections have localized D2-like dopamine receptors to specific neuronal cells in the hippocampus and layers V and VI of the cerebral cortex. This distribution is consistent with known dopaminergic afferent systems to the hippocampus (Goldsmith and Joyce 1994) and to cortical pyramidal neurons of layers V and VI (Goldman-Rakic et al. 1989; Gullledge and Jaffe 1998). Unlike Ntera-2 cells in culture, post-mortem rat brain is exposed to very high levels of released dopamine within a few minutes of death (Vulto et al. 1988) and agonist-induced internalization of dopamine receptors can also be a very rapid process (Vickery and von Zastrow 1999; Macey et al. 2004). It is possible that receptors have been internalized as a result of this process and this may partly explain why their endosomal location has not been described previously. However, the presence of receptors in the cell body, rather than dendritic processes, may make it easier to identify dopamine-responsive neurons by immunohistochemistry. For example, radioligand binding studies detected D2-like receptors in layers V and VI of the cortex, but binding in layers I–III was also found (Martres et al. 1985; Bouthenet et al. 1987). This might be explained by projection of apical dendrites from pyramidal neurons in layers V and VI into layers I–III, since ligand binding detects active receptors on dendrites whereas endosomes will be most evident in the neuronal cell body. These two studies also found D2-like receptors in the hippocampus and this was confirmed by later studies with more specific ligands for DRD2 (Camps et al. 1989) and DRD3 (Levesque et al. 1992; Herroelen et al. 1994; Hall et al. 1996; Bancroft et al. 1998). The distribution of cytoplasmic vesicle labelling is therefore in broad agreement with earlier radioligand studies.

Acknowledgements

This study was supported by a studentship (Robert Hughes Jones Bequest for Research into Schizophrenia) from the Clwyd Institute of Health Studies for E.C.W. We thank Prof. P.W. Andrews (University of Sheffield, UK) for the Ntera-2 cell line. G.E.M. and G.S. thank the British Council in Croatia for support with travel grants.

References

1. Albert PR, Neve KA, Bunzow JR et al (1990) Coupling of a cloned rat dopamine-D2 receptor to inhibition of adenylyl cyclase and prolactin secretion. *J Biol Chem* 265:2098–2104
2. Andrews PW, Damjanov I, Simon D et al (1984) Pluripotent embryonal carcinoma clones derived from the human teratocarcinoma cell line Tera-2. Differentiation in vivo and in vitro. *Lab Invest* 50:147–162
3. Bancroft GN, Morgan KA, Flietstra RJ et al (1998) Binding of [3H]PD 128907, a putatively selective ligand for the D3 dopamine receptor, in rat brain: a receptor binding and quantitative autoradiographic study. *Neuropsychopharmacology* 18:305–316
4. Bouthenet ML, Martres MP, Sales N et al (1987) A detailed mapping of dopamine D-2 receptors in rat central nervous system by autoradiography with [125I]iodosulpride. *Neuroscience* 20:117–155

5. Bunzow JR, Van Tol HH, Grandy DK et al (1988) Cloning and expression of a rat D2 dopamine receptor cDNA. *Nature* 336:783–787
6. Camps M, Cortes R, Gueye B et al (1989) Dopamine receptors in human brain: autoradiographic distribution of D2 sites. *Neuroscience* 28:275–290
7. Defagot MC, Malchiodi EL, Villar MJ et al (1997) Distribution of D4 dopamine receptor in rat brain with sequence-specific antibodies. *Brain Res Mol Brain Res* 45:1–12
8. Dohlman HG, Thorner J, Caron MG et al (1991) Model systems for the study of seven-transmembrane-segment receptors. *Annu Rev Biochem* 60:653–688
9. Ferguson SS (2001) Evolving concepts in G protein-coupled receptor endocytosis: the role in receptor desensitization and signaling. *Pharmacol Rev* 53:1–24
10. Ferguson SS, Downey WE, Colapietro AM et al (1996) Role of beta-arrestin in mediating agonist-promoted G protein-coupled receptor internalization. *Science* 271:363–366
11. Goldman-Rakic PS, Leranth C, Williams SM et al (1989) Dopamine synaptic complex with pyramidal neurons in primate cerebral cortex. *Proc Natl Acad Sci USA* 86:9015–9019
12. Goldsmith SK, Joyce JN (1994) Dopamine D2 receptor expression in hippocampus and parahippocampal cortex of rat, cat, and human in relation to tyrosine hydroxylase-immunoreactive fibers. *Hippocampus* 4:354–373
13. Gullledge AT, Jaffe DB (1998) Dopamine decreases the excitability of layer V pyramidal cells in the rat prefrontal cortex. *J Neurosci* 18:9139–9151
14. Hall H, Halldin C, Dijkstra D et al (1996) Autoradiographic localisation of D3-dopamine receptors in the human brain using the selective D3-dopamine receptor agonist (+)-[3H]PD 128907. *Psychopharmacology (Berl)* 128:240–247
15. Herroelen L, De Backer JP, Wilczak N et al (1994) Autoradiographic distribution of D3-type dopamine receptors in human brain using [3H]7-hydroxy-N,N-di-n-propyl-2-aminotetralin. *Brain Res* 648:222–228
16. Hurlbert MS, Gianani RI, Hutt C et al (1999) Neural transplantation of hNT neurons for Huntington's disease. *Cell Transplant* 8:143–151
17. Levesque D, Diaz J, Pilon C et al (1992) Identification, characterization, and localization of the dopamine D3 receptor in rat brain using 7-[3H]hydroxy-N,N-di-n-propyl-2-aminotetralin. *Proc Natl Acad Sci USA* 89:8155–8159
18. Levey AI, Hersch SM, Rye DB et al (1993) Localization of D1 and D2 dopamine receptors in brain with subtype-specific antibodies. *Proc Natl Acad Sci USA* 90:8861–8865
19. Macey TA, Gurevich VV, Neve KA (2004) Preferential interaction between the dopamine D2 receptor and Arrestin2 in neostriatal neurons. *Mol Pharmacol* 66:1635–1642
20. Martres MP, Bouthenet ML, Sales N et al (1985) Widespread distribution of brain dopamine receptors evidenced with [125I]iodosulpride, a highly selective ligand. *Science* 228:752–755
21. Mauger C, Sivan B, Brockhaus M et al (1998) Development and characterization of antibodies directed against the mouse D4 dopamine receptor. *Eur J Neurosci* 10:529–537

22. Mengod G, Martinez-Mir MI, Vilaro MT et al (1989) Localization of the mRNA for the dopamine D2 receptor in the rat brain by in situ hybridization histochemistry. *Proc Natl Acad Sci USA* 86:8560–8564
23. Monsma FJ Jr, Mahan LC, McVittie LD et al (1990) Molecular cloning and expression of a D1 dopamine receptor linked to adenylyl cyclase activation. *Proc Natl Acad Sci USA* 87:6723–6727
24. Morris GE (1998) Epitope Mapping. In: Rapley R, Walker JM (eds) Humana press, Totowa, NJ, USA, p. 619–630
25. Ng GY, Mouillac B, George SR et al (1994) Desensitization, phosphorylation and palmitoylation of the human dopamine D1receptor. *Eur J Pharmacol* 267:7–19
26. Nguyen TM, Morris GE (1996) Production of panels of monoclonal antibodies by the hybridoma method. *Methods Mol Biol* 66:377–389
27. Oliveri RL, Annesi G, Zappia M et al (2000) The dopamine D2 receptor gene is a susceptibility locus for Parkinson's disease. *Mov Disord* 15:127–131
28. Pereboev AV, Ahmed N, thi Man N et al (2001) Epitopes in the interacting regions of beta-dystroglycan (PPxY motif) and dystrophin (WW domain). *Biochim Biophys Acta* 1527:54–60
29. Prou D, Gu WJ, Le Crom S et al (2001) Intracellular retention of the two isoforms of the D(2) dopamine receptor promotes endoplasmic reticulum disruption. *J Cell Sci* 114:3517–3527
30. Rivera A, Cuellar B, Giron FJ et al (2002) Dopamine D4 receptors are heterogeneously distributed in the striosomes/matrix compartments of the striatum. *J Neurochem* 80:219–229
31. Sodja C, Fang H, Dasgupta T, Ribocco M, Walker PR, Sikorska M (2002) Identification of functional dopamine receptors in human teratocarcinoma NT2 cells. *Brain Res Mol Brain Res* 99:83–91
32. Sokoloff P, Giros B, Martres MP et al (1990) Molecular cloning and characterization of a novel dopamine receptor (D3) as a target for neuroleptics. *Nature* 347:146–151
33. Sorkina T, Hoover BR, Zahniser NR et al (2005) Constitutive and protein kinase C-induced internalization of the dopamine transporter is mediated by a clathrin-dependent mechanism. *Traffic* 6:157–170
34. Sun W, Ginovart N, Ko F et al (2003) In vivo evidence for dopaminemediated internalization of D2-receptors after amphetamine: differential findings with [3H]raclopride versus [3H]spiperone. *Mol Pharmacol* 63:456–462
35. Sunahara RK, Guan HC, O'Dowd BF et al (1991) Cloning of the gene for a human dopamine D5 receptor with higher affinity for dopamine than D1. *Nature* 350:614–619
36. Van Tol HH, Bunzow JR, Guan HC et al (1991) Cloning of the gene for a human dopamine D4 receptor with high affinity for the antipsychotic clozapine. *Nature* 350:610–614
37. Vickery RG, von Zastrow M (1999) Distinct dynamin-dependent and independent mechanisms target structurally homologous dopamine receptors to different endocytic membranes. *J Cell Biol* 144:31–43
38. Vulto AG, Sharp T, Ungerstedt U et al (1988) Rapid postmortem increase in extracellular dopamine in the rat brain as assessed by brain microdialysis. *J Neurochem* 51:746–749

39. Wedzony K, Chocyk A, Mackowiak M et al (2000) Cortical localization of dopamine D4 receptors in the rat brain—immunocytochemical study. *J Physiol Pharmacol* 51:205–221
40. Williams J, Spurlock G, Holmans P et al (1998) A meta-analysis and transmission disequilibrium study of association between the dopamine D3 receptor gene and schizophrenia. *Mol Psychiatry* 3:141–149

Table 1. A panel of monoclonal antibodies against different epitopes of each of the D2-like dopamine receptors

Antibody name	Clone no.	Dopamine receptor	Antigen (amino acids)
MANDOP 31	1D7	DRD3	N-terminus (2–10)
MANDOP 32	3F1	DRD3	N-terminus (2–10)
MANDOP 33	3B7	DRD3	3rd Cytopl. domain (210–218)
MANDOP 34	4D10	DRD3	3rd Cytopl. domain (210–218)
MANDOP 35	9F4	DRD3	3rd Cytopl. domain (210–218)
MANDOP 21	3B9	DRD2	3rd Cytopl. domain (272–282)
MANDOP 22	3D9	DRD2	3rd Cytopl. domain (272–282)
MANDOP 41	2B9	DRD4	3rd Extracell. domain (176–185)

Figure legends

Fig. 1 Monoclonal antibodies against D2-like dopamine receptors recognize proteins of 50 kDa that are associated with membranes. (a) Western blots of human brain cortex extract were developed with three dopamine D3 receptor antibodies at 1/100 dilution of culture supernatant, two mAbs against the N-terminus and one against the 3rd cytoplasmic domain. A control antibody against 43 kDa betadystroglycan [DAG43] (Pereboev et al. 2001) was used as a size marker. (b) Western blots of a different human brain cortex extract developed with all DRD2, DRD3 and DRD4 antibodies at 1/100 dilution of culture supernatant. (c) Subcellular fractionation of Ntera-2 cells by western blotting. Cell pellets were rinsed in PBS, harvested and homogenised in nine volumes of RSB buffer (10mM NaCl, 1.5mM MgCl₂, 10mM Tris-HCl, pH 7.5) using a Dounce homogenizer. The suspension was incubated on ice for 15 min and sonicated before centrifuging at 160,000g for 15 min. The supernatant was removed (cytosol, SN1) and the pellet, which contains all membrane fractions, was resuspended in RSB/1% w/v Triton (P1). The suspension was incubated on ice for 15 min, sonicated and centrifuged as described above. The supernatant (Triton-extractable membrane proteins; SNT) was removed and the pellet was resuspended in RSB buffer (PT). All fractions were boiled with an equal volume of 2x SDS sample buffer [2% SDS, 10% 2-mercaptoethanol, 125 mM Tris-HCl (pH 6.8)] for Western blotting with MANDOP35 mAb against DRD3 (1/100)




Fig. 2 Localization of D2-like dopamine receptors in endosomes in Ntera-2 cells and on dendritic processes in rat brain. All size bars are 10 microns, except where indicated. (a) Each receptor (green indicates FITC conjugated anti-mouse Ig) is localized in cytoplasmic vesicles, often adjacent to the nucleus (red indicates ethidium bromide, 1 lg/ml). The control experiment (inset) shows that DRD3 fluorescence was completely absent when the primary mAb was omitted. (b) DRD3 distribution in Ntera-2 cells was examined by double-labelling with a commercial rabbit polyclonal antibody (red indicates TRITC conjugated anti-rabbit Ig) against the DRD3 N-terminus (Calbiochem) (1/500) and a DRD3 monoclonal antibody (green) against the 3rd cytoplasmic domain (MANDOP 34) (1/4). The overlay shows good colocalization (yellow). (c) Ntera-2 cells were double-labelled with the same rabbit antiserum against DRD3 (1/500: red) and a mouse monoclonal antibody against mannose-6-phosphate receptor [Ab2733 (2G11) Abcam, Cambridge, UK: 1/200: green]. In the overlay, nuclei were counterstained using DAPI (1 µg/ml: blue). (d) Sections from rat basal ganglia were stained with MANDOP34 (1/4) and FITC conjugated anti-mouse (1/50) and counterstained with ethidium bromide (1 lg/ml). DRD3 receptors are localized in thin processes. This staining was absent in controls in which primary mAb was replaced with culture medium (not shown). These photographs were taken with a Zeiss LSM510 Meta confocal system. (e) DRD3 staining of both plasma membrane (left white arrow) and internal vesicles (right white arrow) in Ntera-2 cells. Staining as in (d)




Fig. 3 Distribution of D2-like dopamine receptors in rat cerebral cortex and hippocampus. Five micrometer serial sections were taken through a whole rat brain. The distribution of all three D2-like dopamine receptors in selected sections was examined using DRD2 (MANDOP 21), DRD3 (MANDOP 34) and DRD4 (MANDOP41) (all 1/4) and FITC conjugated anti-mouse Ig (1/50). Cell nuclei were counterstained with ethidium bromide (1 lg/ml). The ethidium bromide stain was photographed at low power (10 \times objective) and a montage of the whole section was produced. High power photographs (63 \times objective) of the regions shown in boxes were used to determine the subcellular distribution of D2-like dopamine receptors. (a) Cerebral cortex. The boxes labelled 1–6 correspond to cortical layers I–VI. Layers III and IV, like layer II, showed no antibody staining for receptors (not shown). The “control + antigen” inset illustrates the prevention of antibody staining by pre-incubation with peptide immunogen; in this case, MANDOP34 (left, green) on layer V cells in the cortex; ethidium bromide counterstain (right, red). (b) Hippocampus. The montage includes granular neurons of the dentate gyrus and CA1, CA2 and CA3 neurons of the hippocampus. CA3 neurons showed consistent staining of cytoplasmic vesicles for all three receptors, as did occasional neurons in the polymorphic cell layer and within the stratum radiatum and lacunosum regions of the hippocampal formation

